



# Low birth weight

## What it is and why it matters

Low birth weight refers to babies that weigh less than 5-½ pounds when they're born. Low-birth-weight babies are more likely to get sick — or die — in the first year.

## Tips to help prevent low birth weight

A healthy pregnancy starts with making healthy lifestyle choices like:



**Go to all your prenatal care visits** – Prenatal care is the care a woman gets during pregnancy. Regular checkups let your provider check the health of you and your baby.



**Get tested for sexually transmitted diseases (STDs)** – STDs can be harmful for pregnant women and their babies. Talk to your provider about STD tests available. See the back side to learn more.



**Quit smoking** – Smoking can lead to low birth weight, miscarriage and other problems. MI Tobacco Quitlink offers free help at 1-800-QUIT-NOW (1-800-784-8669) or [michigan.quitlogix.org](http://michigan.quitlogix.org).



**Eat a healthy diet** – Eat fresh fruits and vegetables if you can. Drinking plenty of water and taking vitamins during pregnancy are important, too.

## Get extra support

Build a healthy future and earn great rewards with Healthy First Steps®. This program will help you make your pregnancy the healthiest it can be and give your baby the best start possible.

To sign up, call toll-free **1-800-599-5985**, TTY **711**, 8 a.m.–5 p.m., Monday–Friday.

See other side to learn more →

## Get tested for STDs throughout your pregnancy

It's important to get these tests as part of your regular prenatal checkups:

- **First trimester visit** – syphilis, HIV/AIDS, hepatitis B and hepatitis C, chlamydia and gonorrhea
- **Third trimester visit** – for those at risk of infection or reinfection: syphilis, HIV/AIDS, chlamydia, gonorrhea and hepatitis C
- **At delivery** – for those who haven't yet been tested or are at risk of reinfection: syphilis, HIV/AIDS, hepatitis B and hepatitis C

### Know the danger of congenital syphilis

This disease happens when a woman with syphilis passes it on to her baby during pregnancy. There are often no symptoms. But it can be easily cured with antibiotics.

Congenital syphilis can cause:

- Miscarriage or still birth
- Premature (early) birth
- Low birth weight
- Death shortly after birth

[myuhc.com/communityplan](https://myuhc.com/communityplan)



UnitedHealthcare does not discriminate on the basis of race, color, national origin, sex, age or disability in health programs and activities.

We provide free services to help you communicate with us. Such as, letters in other languages or large print. Or, you can ask for an interpreter. To ask for help, please call the toll-free member phone number listed on your health plan member ID card, TTY 711, 8:30 a.m.–5:30 p.m., Monday–Friday.

ATENCIÓN: si habla español, hay servicios de asistencia de idiomas, sin cargo, a su disposición. Llame al número de teléfono gratuito que aparece en su tarjeta de identificación.

مجاناً. اتصل على الرقم المجاني المدون على بطاقة هويتك تنبيه: إذا كنت تتحدث العربية، فتتوفر خدمات المساعدة اللغوية